

SOME MORPHOLOGICAL TRAITS OF THE TARSUS FORK-NOSE DOG IN TURKEY

YILMAZ, Orhan

Igdir University, Faculty of Agriculture, Department of Animal Science, 76100, Igdir.

ABSTRACT

This research is the first to document the morphology of the Tarsus Fork-nose Dog in Turkey. The purpose was to determine the morphological traits of this dog in different age and sex groups as raised in the South of Turkey. To this end, a total of 24 male and 31 female Fork-nose were measured. Descriptive statistics gave the following means: live weight was 21.7 kg, withers height 48.5 cm, height at rump 48.5 cm, body length 49.1 cm, heart girth circumference 64.0 cm, chest depth 20.8 cm, chest width 18.8 cm, cannon circumference 10.5 cm, and head length 20.7 cm respectively.

Keywords: Tarsus fork-nose, morphological trait, body measurements, coat colour.

Correspondence: zileliorhan62@hotmail.com

Office: +90-4762261314/1225 Fax: +90-4762261139

INTRODUCTION

The Tarsus fork-nose dogs have a fully split nose, essentially resulting in a double nose, that is quite a rare feature in dog breeds. 'Fork-nose' describes a nose split vertically right down the middle, accounting for these particular dogs being described as 'fork-nose hunters' (www.thebreedsofdogs.com 2011). Something approaching a split appears to be normal for the nose of a dog. In most dogs, the nostrils are separated vertically by a band of skin and fur dividing the nose all the way to the dog's upper lips. Even though there are no researches to indicate whether the fully-split double nose is a benefit or a hindrance, many people believe that the split nose increases the scent discrimination ability of a dog possessing it (en.wikipedia.org 2011)



Figure 1. An adult Fork-nose dog.

The Turkish sighthound, of which the Tarsus Fork-nose dogs are a subgroup, is generally located in the Province of Icel, Mediterranean region (Yilmaz 2007, Yilmaz 2008,). Thus the Tarsus Fork-nose is a dog that has been bred primarily for hunting, and it hunts by sight as well as scent. It is, however, particularly good at hunting by scent and indicating the prey as a pointer dog. These

dogs have a very acute sense of smell, possibly heightened by their peculiar nose structure. This helps them greatly when they are hunting, conferring superior tracking abilities. Their sense of smell is thus what makes them excellent hunting dogs. At the same time they are very friendly dogs, getting along with humans as well as other dogs.



Figure 2. An adult Fork-nose dog.

Tarsus Fork-nose dogs are not very furry. They have short-haired coats, usually seen in brown and cream that are very easy to maintain. Some of the brown dogs are pied, that is a combination of brown and white. Although they are quite affectionate dogs, Fork-nose dogs are not suited for apartment life and need lots of space move around (Yılmaz 2007, Yılmaz 2008).



Figure 3. A Fork-nose puppy.

Apart from the Tarsus Fork-nose, there are several fork-nose dog breeds elsewhere in the world including the Pachon Navarro (Old Spanish Pointer) in Spain and the Andean Tiger Hound in Bolivia (www.thebreedsofdogs.com 2011, en.wikipedia.org 2011).

The Pachon Navarro is a native Braque hunting dog of Spain. They have been known since the 1800s and appear in a number of oil paintings. The Pachon Navarro can either be long haired or short haired. They are described as very friendly, loyal and brave hunters. These dogs, too, are characterized by their ability to point in the direction of the game. This ability naturally makes them great hunting companions. The Navarro also has immense physical strength and can fight other bigger animals, but can be used as pet dogs in the home as well because of their sunny disposition. Navarro dogs are very hard to ignore because of their substantial size: 20-33 kg in weight and, 45 to 60 cm in height average, seen in Table 1 (www.sobreperros.com 2011, www.dogbreeds.net 2011, www.pachonnavarro.com 2011, www.continentalkennelclub.com 2011, www.pachon.info 2011).

Table 1. Weight and height of Pachon Navarro Dogs.

Source	Weight (kg)			Height (cm)		
	General	Male	Female	General	Male	Female
www.thebreedsofdogs.com ¹	27-33			48-57		
en.wikipedia.org ²	27-33			48-57		
www.sobreperros.com ⁵	27-33				55-64	53-60
www.dogbreeds.net ⁶	27-33			48-59		
www.pachonnavarro.com ⁷		28	25	45-60	55	52
www.continentalkennelclub.com ⁸	25-30				51-61	
www.pachon.info ⁹	20-30				55	52

The Double-nosed Andean Tiger Hound is another of the rare split-nose breeds in the World, and lives in Bolivia. It is believed that the Andean Tiger is descended from the Pachon Navarro, assuming it to have been brought to Bolivia by the Spanish Conquistadors in the 16th century (en.wikipedia.org 2011).

The object of this study was to determine body coat colour and some phenotypic characteristics of the Tarsus Fork-nose dogs.

MATERIALS AND METHODS

Experimental animals

The Fork-nose dogs in the study were surveyed in January 2011 in the city of Tarsus, in the Province of Icel, in the South of Turkey (36^o 55'N; 34^o 54'E) (www.googleearth.com). A total of 55 dogs, 24 male and 31 female, were studied. The sampled dogs were aged between 1 and 10 years, and divided into four age groups: 1-2 years, 3-4 years, 5-6 years and 7-10 years. In the first group there were 4 males and 5 females; in the second group there were 12 males and 9 females; in the third group there were 4 males and 13 females; and in the fourth group there were 4 males and 4 females. Almost all dogs belonged to registered hunters, and the ages of dogs were determined from their ID cards.

Measurements

The sampled dogs were weighed for live weight (LW) with a portable spring scale. Linear measures such as withers height (WH), height at rump (HR), body length (BL), chest depth (CD), and chest width (CW) were measured using a measuring stick calibrated in centimetres. Other linear measures such as heart girth circumference (HGC), cannon circumference (CC), and head length (HL) were measured using a graduated plastic tape (Yılmaz 2007).

Statistical analysis

The data obtained were analyzed with the Minitab 15 statistical software program. Descriptive statistics for body dimensions were analyzed using ANOVA and Student's T-Test that also

determined the impact of sex, region, body coat colour, and age group on the response variables of LW, WH, HR, BL, HGC, CD, CW, CC, and HL (Anonymous, 2011).

RESULTS AND DISCUSSION

Descriptive statistics for the morphological traits are given in Table 2. Regarding to the effect of sex on phenotypic traits, analyzed t-Test results show that there are significant differences between male and female dogs for LW, HR, BL, CD (P<0.01) and WH and HGC (P<0.05). For all results significant or not, male dogs yielded higher values than females.

The age means are given in Table 2. In the 1-2 years age group Fork-nose dogs were significantly different from the other age groups for all traits (P<0.05). After the age of 2 years there were minor differences for live weight and body dimensions. It can be concluded that Tarsus Fork-nose dogs continue growing up to 2 years of age, and then their growth rate slows.

Table 2. Descriptive statistics and comparison results of the phenotypic traits of Tarsus Fork-nose for different sexes and ages.

Trait		LW (kg)	WH (cm)	HR (cm)	BL (cm)	HGC (cm)	CD (cm)	CW (cm)	CC (cm)	HL (cm)
		$\bar{X} \pm S_{\bar{X}}$								
Sex	Overall (n=55)	21.7 ± 0.53	48.5 ± 0.69	48.5 ± 0.65	49.1 ± 0.81	64.0 ± 0.69	20.8 ± 0.48	18.8 ± 0.34	10.5 ± 0.17	20.7 ± 0.30
	Male (n=24)	23.6b** ± 0.93	50.3b* ± 1.09	50.5b** ± 1.04	51.5b** ± 1.01	65.7b* ± 0.75	22.4b** ± 0.58	19.0a ± 0.37	10.7a ± 0.23	20.8a ± 0.50
	Female (n=31)	20.3a** ± 0.49	46.9a* ± 0.79	46.9a** ± 0.73	47.2a** ± 1.09	62.7a* ± 1.03	19.6a** ± 0.64	18.6a ± 0.53	10.4a ± 0.23	20.5a ± 0.50
Age	1-2 years (n=9)	12.8a** ± 0.31	41.6a** ± 0.99	42.8a** ± 1.14	42.2a** ± 1.74	59.1a** ± 2.38	17.7a** ± 1.18	16.7a** ± 1.18	8.9a** ± 0.29	17.6a** ± 0.63
	3-4 years (n=21)	17.0b** ± 0.63	49.3b** ± 1.10	49.5b** ± 1.09	50.2b** ± 1.17	64.6b** ± 0.86	21.1b** ± 0.65	18.5ab** ± 0.42	10.5a** ± 0.18	21.4b** ± 0.37
	5-6 years (n=17)	16.5b** ± 0.50	49.5b** ± 0.89	49.1b** ± 0.91	50.4b** ± 1.24	64.5b** ± 0.99	21.0b** ± 0.90	19.4b** ± 0.42	11.1b** ± 0.23	20.9b** ± 0.40
	7-10 years (n=8)	18.3b** ± 1.21	51.4b** ± 1.22	50.9b** ± 1.25	51.1b** ± 1.93	66.6b** ± 1.71	23.1b** ± 0.90	20.6b** ± 0.22	11.3b** ± 0.50	21.6b** ± 0.71

* (P<0.05), ** (P<0.01),

There were no significant differences between means showed in same letters within a column and factor group

Table 3 shows the coat colour distributions of Fork-nose dogs. The Tarsus Fork-nose dogs, nearly 40% are pied, white-brown and white-black mixes. In total more than 50% of the dogs are in brown colour.

Table 3. Distributions of body coat colour of Tarsus Fork-nose.

Trait	White and Brown	White and Black	Brown	Black
n	13	8	29	5
%	23.6	14.6	52.7	9.1

The phenotypic correlation values displayed in the Table 4 show that most of observed traits are affected by selected factors. The highest values was found between WH and HR ($r = 0.98$) ($P < 0.01$). Other high values were found between LW and WH ($r = 0.69$), LW and HR ($r = 0.68$), WH and CD ($r = 0.69$), HGC and CW ($r = 0.69$) ($P < 0.01$). The correlations of LW-HL, WH-BL, WH-CC, HR-BL, HR-CD, and BL-CC also yielded higher values those than $r = 0.60$ ($P < 0.01$). The lowest value ($r = 0.28$) was found between BL and HL ($P < 0.05$). Other low correlation values were found between WH and AD ($r = 0.34$), HR and CW ($r = 0.33$), and CD and CC ($r = 0.33$) ($P < 0.05$). There were no negative correlations between all other traits, as seen in Table 4.

Table 4. Phenotypical correlation coefficients (r) between body measurements in Fork-nose dogs.

Traits	LW	WH	HR	BL	HGC	CD	CW	CC
WH	0.69**							
HR	0.68**	0.98**						
BL	0.46**	0.63**	0.63**					
HGC	0.50**	0.48**	0.50**	0.55**				
CD	0.39**	0.69**	0.64**	0.19	0.25			
CW	0.25	0.34*	0.33*	0.49**	0.69**	0.23		
CC	0.58**	0.62**	0.59**	0.60**	0.55**	0.33*	0.35**	
HL	0.62**	0.57**	0.58**	0.28*	0.45**	0.25	0.16	0.49**

* $P < 0.05$, ** $P < 0.01$

As yet there has been no literature report revealing the morphological features of the Tarsus Fork-nose dog, so from overall results of the current study, we may conclude that Tarsus Fork-nose Dog is small size hunting dog. Although brown and white-brown coat colours are predominant, Tarsus Fork-nose Dog owners make no selection on body coat colour. The dogs grow up to 2 years of age and that there is only minor growth. This suggests that Tarsus Fork-nose Dog reaches mature body weight and size at around 2 years of age. In order to establish whether or not they are descended from one ancestor, a full DNA based research can be conducted in the future, and a genetic relationship sought between the Tarsus Fork-nose, Pachon Navarro and the Double-nosed Andean Tiger dogs. With regard to the Tarsus Fork-nose, a larger sample dogs can be used to research of dog behaviour and hunting performance.

BIBLIOGRAPHY

1. Anonymous, 2011. Minitab 15 Computer Program.
2. Yılmaz O: Some Morphological Characteristics of Kangal Dogs Raised in Various Regions of Turkey (unpublished PhD study). pp. 50-59, University of Ankara, Faculty of Agriculture, Department of Animal Science, Ankara, 2007.
3. Yılmaz O: Turkish Kangal (Karabash) Shepherd Dog. First ed., pp. 24, Impress Printing, Ankara, 2008.
4. en.wikipedia.org/wiki/Double-nosed_Andean_tiger_hound (accessed on 14.03.2011)
5. www.continentalkennelclub.com (accessed on 14.03.2011)
6. www.dogbreeds.net/pachon-navarro (accessed on 14.03.2011)
7. www.googleearth.com (accessed on 28.02.2011)
8. www.pachonnavarro.com/pdf/Estandar (accessed on 14.03.2011)
9. www.pachon.info/contenido (accessed on 13.02.2011)
10. www.sobreperros.com/raza/pachon_navarro (accessed on 14.03.2011)
11. www.thebreedsofdogs.com/pachon-navarro (accessed on 14.03.2011)